CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM

maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

and supported by the

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

Database Version 8.1 (2005)

B123 Red-tailed Hawk Buteo jamaicensis

Family: Accipitridae Order: Falconiformes Class: Aves

Written by: C. Polite, J. Pratt Reviewed by: S. Bailey Edited by: S. Bailey

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

Common, permanent breeding and winter resident and migrant. Breeds throughout California, and winters in all areas without heavy snow cover. Found in almost all habitats, from lowest to highest elevations.

SPECIFIC HABITAT REQUIREMENTS

Feeding: Eats small mammals up to hares in size, small birds, reptiles, amphibians, and some carrion (Orians and Kuhlman 1956). In winter, largely dependent upon mice, but also takes medium to fairly large birds on the ground. Catches small birds during migration. Searches by soaring; also perches and pounces, or pounces on prey from low, quartering flights, sometimes hovering on wind or air currents.

Cover: Feeds in grasslands and grass/shrub stages of most habitats. Roosts in trees; sometimes in dense conifer stands.

Reproduction: Usually nests in large trees near openings, in older, mature forests, especially riparian deciduous habitats. Occasionally nests on cliffs or low ledges. Nests 9-21 m (30-70 ft) above ground in trees, higher on cliffs. Flexible in choice of nest site; occasionally uses human-made structures, shrubs, cacti. Sometimes nests in isolated trees, or in small groves in open habitat.

Water: Requirements probably met from food.

Pattern: Highly adaptable; uses grasslands, open brush habitats, and open stands of deciduous and conifer forests. Also frequents croplands, fields, and pastures.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity. Most active feeding occurs in early morning and late afternoon.

Seasonal Movements/Migration: Migrates downslope in winter; in summer and fall, some individuals move to open areas at higher elevations.

Home Range: Home ranges vary from less than 1 to 10 km² (0.3 to 3.8 mi²), depending on location, topography, habitat, and prey availability.

Territory: In California, Fitch et al. (1946) calculated that territories varied from 0.3 to 0.8 km² (0.1 to 0.3 mi²). They found 0.8 breeding pairs/km² (2/mi²). Territory defended yearlong.

Reproduction: Courtship begins as early as January. Breeds March through July; peaking in May and June. Clutch of 2-5 eggs, usually 2-3, laid in March and April. Incubates 28-32 days. Semialtricial young fledge in 40-45 days.

Niche: Adaptable, common, and widespread. Scrub jays, and other avian and mammalian predators, take undefended eggs and nestlings (Brown and Amadon 1968). May compete for food with ferruginous, Swainson's, and rough-legged hawks. Great horned owls commonly use old nests, and occasionally golden eagles do. Nestlings may be killed by blood-sucking flies (Fitch et al. 1946). Golden eagles may prey on adult.

REFERENCES

- Brown, L., and D. Amadon. 1968. Eagles, hawks and falcons of the world. 2 Vols. Country Life Books, London. 945pp.
- Call, M. W. 1978. Nesting habits and survey techniques for common western raptors. U. S. Dep. Inter., Bur. Land Manage., Portland, OR. Tech. Note No. 316. 115pp.
- Craighead, J. J., and F. C. Craighead, Jr. 1956. Hawks, owls and wildlife. Stackpole Books, Harrisburg, PA. 443pp.
- Fitch, H. S., R. Swenson, and D. F. Tillotson. 1946. Behavior and food habits of the redtailed hawk. Condor 48:205-237.
- Gates, J. M. 1972. Red-tailed hawk populations and ecology in east-central Wisconsin. Wilson Bull. 84:421-433.
- Jackman, S. M., and J. M. Scott. 1975. Literature review of twenty three selected forest birds of the Pacific Northwest. U.S. Dep. Agric., For. Serv., Reg. 6, Portland OR. 382pp.
- Luttich, S. N., L. B. Keith, and J. D. Stephenson. 1971. Population dynamics of the red-tailed hawk (Buteo jamaicensis) at Rochester, Alberta. Auk 88:75-87.
- Maser, C., B. R. Mate, J. F. Franklin, and C. T. Dyrness. 1981. Natural history of Oregon coast mammals. U.S. Dep. Agric., Pac. Northwest For. and Range Exper. Sta., Portland. Gen. Tech. Rep. PNW-133. 496pp.
- Orians, G. H., and F. Kuhlman. 1956. Red-tailed hawk and horned owl populations in Wisconsin. Condor 58:371-385.
- Wiley, J. W. 1975a. The nesting and reproductive success of red-tailed hawks and red-shouldered hawks in Orange County, California, 1973. Condor 77:133-139.
- Wiley, J. W. 1975b. Relationships of nesting hawks with great horned owl. Auk 92:157-159.